

### Synonym

FLJ18683,T3E,TCRE,CD3E,CD3-epsilon

### Source

MABSol® Biotinylated Human CD3E, His Tag, primary amine labeling (CDE-H8223) is expressed from human HEK293 cells. It contains AA Asp 23 - Asp 126 (Accession # [NP\\_000724.1](#)). It is the biotinylated form of Human CD3 epsilon Protein (Cat # CDE-H5223).

Predicted N-terminus: Asp 23

### Molecular Characterization

CD3E(Asp 23 - Asp 126)  
NP\_000724.1 Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 12.6 kDa. The protein migrates as 15-16 kDa under reducing (R) condition (SDS-PAGE).

### Biotinylation

*The primary amines in the side chains of lysine residues and the N-terminus of the protein are conjugated with biotins using standard chemical labeling method. A standard biotin reagent (13.5 angstroms) is used in this product.*

### Biotin:Protein Ratio

The biotin to protein ratio is 2-3 as determined by the HABA assay.

### Endotoxin

Less than 1.0 EU per µg by the LAL method.

### Purity

>90% as determined by reduced SDS-PAGE.

### Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 12 months under sterile conditions after reconstitution.

### SDS-PAGE




Biotinylated Human CD3E, His Tag, primary amine labeling on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.


### Bioactivity-ELISA

**Biotinylated Human CD3E, His Tag, primary amine labeling ELISA**

0.2 µg of Mouse Anti-Human CD3 (SP34-2) per well



**Batch consistency**



Lot. No.	EC50(µg/mL)
Lot. No. 1	0.0018
Lot. No. 2	0.0019
Lot. No. 3	0.0017

[Report](#)

Immobilized Mouse Anti-Human CD3 (SP34-2) at 2 µg/mL (100 µL/well) can bind Biotinylated Human CD3E, His Tag, primary amine labeling (Cat. No. [CDE-H8223](#)) with a linear range of 0.1-2 ng/mL (QC tested).

**Background**

CD3e molecule, epsilon is also known as CD3E, is a T-cell surface single-pass type I membrane glycoprotein. CD3E contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. CD3E, together with CD3-gamma, CD3-delta and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. CD3E plays an essential role in T-cell development, and defects in CD3E gene cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to type I diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.

**References**

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.